



SYNTHESIS AND BIOLOGICAL EVALUATION OF NOVEL 3,5-DISUBSTITUTED 1,2-ISOXAZOLINYL DERIVATIVES OF NAPHTHALENE-1-AMINE

*,¹Jawahar M, ²Arunkumar S, ¹Thangamuthu S, ¹Vasavi G, ¹Hemalatha CH

^{*,1}Smt.Sarojini Ramulamma College of Pharmacy, Mahabubnagar, AP, India - 509 001.

²SRM College of Pharmacy, Kattankulathur, Kanchipuram District, TN, India - 603 203.

Abstract

Naphthalene-1-amine known as 1-amino naphthalene is an aromatic amine possesses selective serotonin reuptake inhibitor and antimicrobiological activity. New isoxazoline derivative was synthesized from α -naphthylamine to get better biological activities like anti cancer, antibiotic, antifungal, antiviral, anti-inflammatory, antidepressant, anthelmintic. The synthesized novel heterocyclic derivative i.e isoxazoline compounds or derivatives were screened for their antimicrobial and anticancer activity and compared with standard drugs.

Keywords: Heterocyclic derivatives, Isoxazoline, Chalcone, Antifungal, Anticancer, MTT Assay.

Introduction

Synthesis of α -naphthylamine on Acetylation reaction gives substituted aromatic ketone . The base catalyzed involves condensation of substituted aromatic ketones and substituted aldehydes to give α,β -unsaturated ketones (Chalcones), which on cyclization with hydroxylamine hydrochloride in alkaline medium gave corresponding Isoxazoline derivatives⁷. Literature review reveals that isoxazoline derivatives exhibit pharmacological activities such as Anti-inflammatory, Antitubercular, Antibacterial, Antifungal, Antiviral, Analgesic, Antitumour, Anticoagulant, Antidepressant, Antipsychotic etc. Based on above facts it is worthwhile to

prepare Isoxazolinyl derivatives of Naphthalene-1-Amine¹⁻⁹ (α -Naphthylamine).


Experimental procedure synthesis

Synthesis of N – (naphthalene – 1- yl) acetamide

To a solution of naphthalen -1 – amine¹ (0.01 mole) in chloroform (dry, 100ml), acetyl chloride (0.02 mole) is added drop wise at 0-5°C with constant stirring. The reaction mixture was stirred for 2hrs by magnetic stirrer. The excess solvent was distilled off and the separated mass was poured into ice water and recrystallized from methanol.

Author for Correspondence:

Jawahar M,
 Smt.Sarojini Ramulamma College of Pharmacy,
 Mahabubnagar, AP,
 India - 509 001.
 Email: jawapharma@yahoo.com

Scheme 1:**Synthesis of N-(naphthalene – 1- yl) -3- aryl acrylamide derivatives (R₁-R₁₀)**

To a mixture of N- (naphthalene-1-yl) acetamide (0.01 mole) in methanol (50 ml), appropriate aromatic aldehyde¹⁻⁹. (0.01 mole) are added in the presence of 2% NaOH solution (5ml). The reaction mixture is stirred for 10 hrs at room temperature and

then refluxed for 6hrs. The excess solvent was distilled off and poured into ice water. The resulting solid thus separated, is filtered, washed with water and recrystallized from ethanol.

Synthesis of N – (naphthalene – 1- yl) – 5-phenyl - 4, 5 - dihydroisoxazol- 3- amine derivatives (S₁–S₁₀)

To a mixture of N – (naphthalene -1-yl) -3-aryl acryl amide derivative (0.01mole) in absolute ethanol (50, dry), hydroxylamine hydro chloride (0.01 mole) and solid NaOH (0.4g) were added. The reaction mixture was refluxed for 5 hrs and poured in to ice water. The solid thus separated was filtered, washed with water and recrystallized from acetone. The recrystallized products purity was checked by TLC by using solvent system Ethanol, Dichloromethane (1:2) ratio.

Experimental

Physicochemical Parameters

Physical characterization of synthesized compounds was characterized. Melting points were measured in open-end capillary tube method by VEEGO digital electrically heating melting point apparatus. The purity of new compounds and reaction completion was checked by TLC by measuring R_F value of products. Solubility was determined by using various organic solvents like DMSO, Chloroform and Ethanol. Physico-chemical properties are recorded in Table no. 1.

Table No. 1: Physico-Chemical Parameters

Compound code	Solubility	Color	Percentage yield	Melting point	Molecular weight	Molecular formula	Rf value
S ₁	Ethanol, Chloroform, DMSO	Pale violet	85	90-97	304.34	C ₁₉ H ₁₆ N ₂ O ₂	0.95
S ₂	Ethanol, Chloroform, DMSO	Reddish brown	82	110-113	322.78	C ₁₉ H ₁₅ Cl N ₂ O	0.83
S ₃	Ethanol, Chloroform, DMSO	Grey	80	128-130	288.34	C ₁₉ H ₁₆ N ₂ O	0.78
S ₄	Ethanol, Chloroform, DMSO	Grey	80	105-108	348.39	C ₂₁ H ₂₀ N ₂ O ₃	0.89
S ₅	Ethanol, Chloroform, DMSO	Violet	73	128-130	304.34	C ₁₉ H ₁₆ N ₂ O ₂	0.97
S ₆	Ethanol, Chloroform, DMSO	Grey	78	94-95	333.34	C ₁₉ H ₁₅ N ₃ O ₃	0.84
S ₇	Ethanol, Chloroform, DMSO	Pale pink	93	115-120	334.51	C ₂₀ H ₁₉ N ₂ O ₃	0.94
S ₈	Ethanol, Chloroform, DMSO	Pink	75	107-110	322.78	C ₁₉ H ₁₅ Cl N ₂ O	0.86
S ₉	Ethanol, Chloroform, DMSO	Yellow	83	117-120	333.34	C ₁₉ H ₁₅ N ₃ O ₃	0.75
S ₁₀	Ethanol, Chloroform, DMSO	Purple	84	107-110	348.43	C ₂₁ H ₂₀ N ₂ O ₃	0.93

Biological Evaluation

Anti-bacterial activity

Antibacterial activity of the synthesized compound was determined using Serial dilution method. Nutrient broth was used as culture media for the study. Various gram positive and gram negative bacteria were used. Standard antibiotic as Ciprofloxacin was used for comparison of activity. Nutrient broth solution was sterilized by autoclaving at 15lbs pressure for 20 min. Inoculation of bacterial strains were made in above media and incubated at 37°C for 24 hrs. Uninoculated sterile media was used as blank. Test compound was dissolved in DMSO and dilution was made to get desired concentration of 100µg/ml, 50µg/ml, 25µg/ml and 12.5µg/ml. The test tubes were visualized under bright light. The test tube with least concentration of compound showing no bacterial growth was taken as minimum inhibitory concentration (MIC) of the compound.

Anti-fungal activity

Asperigillus niger was used for the Serial dilution study. Sabouraud dextrose broth

cultured with fungal strain was taken in series of test tubes. Test compound was dissolved in DMSO of two-fold decreasing concentrations, incubated at 37° c for 48 hrs. Uninoculated sterile media was used as blank. Ketaconazole was used as standard. The test tubes were under bright light. Test tube with least concentration of compound showing no fungal growth was taken as (MIC) of the compound.

Evaluation of anti-cancer activity

Microculture tetrazolium assay (MTT) was used for evaluating anticancer activity. All the synthesized compounds were evaluated for cytotoxicity using MTT assay. It is a standard colorimetric assay, which measures changes in colour for the determination of viable cells. Assay is dependent on the activity of mitochondrial dehydrogenase enzymes that reduce yellow 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide(MTT) to a blue colour formazoin product. The activity of the enzyme is directly proportional to cell viability. Doxorubicin was used as standard.

$$\text{Cell viability (\%)} = \frac{\text{Mean od} \times 100}{\text{Control of o.d.}}$$

Table No. 2: Antimicrobial and Anticancer activity of synthesized compounds

Compound (100 µg/ml – 12.5µg/ml)	Minimum inhibitor concentration(µg/ml)				IC ₅₀ (mg/ml)
	Antibacterial activity			Antifungal activity	Anticancer activity
	E.coli	S.aureus	B.subtilis	A.niger	HEP2 Cells
S ₁	50	25	50	25	0.056
S ₂	100	50	25	50	0,127
S ₃	25	100	50	50	0.059
S ₄	50	50	50	25	0.133
S ₅	50	25	100	100	0.127
S ₆	25	100	25	25	0.129
S ₇	100	50	50	100	0.111
S ₈	25	25	25	25	0.131
S ₉	50	100	50	100	0.065
S ₁₀	100	50	100	50	0.131
Standard	12.5	12.5	12.5	12.5	0.002

Result and Discussion

All the synthesized compounds were characterized by TLC, Melting point and Solubility. The compounds were evaluated for their antimicrobial activity i.e antibacterial and antifungal, and anticancer activity. Minimum inhibitory concentration value of the compounds (S₁, S₂, S₃, S₅, S₆, S₈) against the bacterial, fungal strain showed moderate inhibition at 25 µg/ml concentration. The bacterial screening indicated that the test compounds were found to be less active, when compared to that of standard drug Ciprofloxacin. The fungal screening indicated that the test compounds, were found to be moderately active, when compared to that of standard drug Ketaconazole. Anticancer activity evaluated by Micro culture tetrazolium assay (MTT) (S₁, S₃, S₇, S₉) In HEP2 cell lines, were found to have moderate activity compared to standard.

Conclusion

The synthesized compounds subjected to antimicrobial and anticancer activity showed moderate activity when compared with standard drug.

References

1. Agarwal.O.P, Organic Chemistry Reactions and Reagents 42th edition 234.
2. Wolff, M.E, Burger Medicinal Chemistry and Drug Discovery, 5th edition, A-Wiley-Inter Science Publication 1997, 2292 - 2296.
3. Block J.H, Beale JM Jr. Text Book of Organic Medicinal and Pharmaceutical Chemistry. Wilson and Gisvold's 11th edition, Lippincott Williams & Wikis Publication, 2004; 01-02.
4. Finar.I.L Organic Chemistry 6th edition; 1994, 803.
5. Surendra N Pandeya. A Text Book of Medicinal Chemistry, 3rd edition, SG Publisher, Varanasi, 2004; 02:1038-1040.
6. David A-William, Thomas L-Lemke Foye's Principles of Medicinal Chemistry, 5th edition, Lippincott Williams and Wilkins, 2002:758-790.
7. Lokanathan Rai, Alfred Hassner. Intermolecular 1, 3-dipolar Cyclo addition of Nitrile Oxides with Vinyl Acetate and Acrylonitrile. Indian Journal of Chemistry. 1997. 36B: 242-245.
8. Lokanatha Rai, .Jaya Shankara.B, Synthesis and Antimicrobial Studies of a New Series of bis-hetero Cycles. European Journal of Chemistry. 2008; 5:2: 370-376.
9. Ashok kumar, Preeti Rani, Srivastava VK. Synthesis and Anti inflammatory activity of Isoxazoliny derivatives of anthranilic acid, Indian Journal of Chemistry. 2003; 42B: 1729- 1733.